



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

BIBLIOGRAPHIC RECORD TARGET

Graduate Library
University of Michigan

Preservation Office

Storage Number: _____

ABN2660

UL FMT B RT a BL m T/C DT 07/18/88 R/DT 07/18/88 CC STAT mm E/L 1

010: : |a 03019635

035/1: : |a (RLIN)MIUG86-B42083

035/2: : |a (CaOTULAS)160031836

040: : |a MiU |c MiU

050/1:0 : |a QA467 |b .W18

100:1 : |a Wallace, Ebenezer Erskine.

245:04: |a The circle squared; |b a solution of the problem of squaring the circle, showing the exact ratio between radius and quadrant, |c by Ebenezer Erskine Wallace.

260: : |a Monmouth, Ill., |b E.E. Wallace, |c 1903.

300/1: : |a 16 p. |b diagrs. |c 23 cm.

650/1: 0: |a Circle-squaring

998: : |c DPJ |s 9124

Scanned by Imagenes Digitales
Nogales, AZ

On behalf of
Preservation Division
The University of Michigan Libraries

Date work Began: _____
Camera Operator: _____

THE CIRCLE SQUARED

A SOLUTION

OF THE

PROBLEM OF SQUARING THE CIRCLE

Showing the Exact Ratio between

RADIUS AND QUADRANT

BY

Ebenezer Erskine Wallace

E. E. WALLACE
MONMOUTH, ILLINOIS
1903

Copyright, 1903,
BY
EBENEZER ERSKINE WALLACE

PREFACE

Up to the present time the problem of finding the exact ratio between the diameter and the circumference of a circle has remained unsolved. For more than one hundred years the leading scientific societies of the world have declared the problem to be impossible of solution. It is, therefore, with some diffidence that I submit the following pages to the consideration of mathematicians and others interested in the solution of a problem heretofore deemed unsolvable. And yet, having worked on the problem for a period of time covering more than fifteen years, it is with considerable assurance of its correctness that I submit the solution, as I believe it to be.

I may be pardoned for such assurance when it is considered that I have worked out the problem by two separate and distinct theorems and find the answers to be the same to six decimal places. This would be impossible if either theorem were incorrect. The answer thus obtained for the ratio of the circumference to the diameter to six decimal places is $3.141941+$. By calculating upon the old theory of approximation and subdividing the sides until I had over three million sides to a polygon inside of a circle, I obtained the answer $3.1418+$, which tends to show that the theorems herewith submitted are correct.

I am well aware that the correctness of these theorems will be called in question, and if they are not mathematically correct they must fall. I submit them, however, with the confident belief that the time will come when they will be universally recognized as the proper solution of this problem.

EBENEZER ERSKINE WALLACE.

MONMOUTH, ILLINOIS, May 18, 1903.

TABLE A

Sides			Remainders or Loss	
6 ×	0	.5	-----	= 3.
12 ×	1	.258819045102520	8460462834	= 3.105828541230240
24 ×	2	.130526183513181	3899039747	= 3.132628404316344
48 ×	3	.065403124924338	0515367387	= 3.139349996368224
96 ×	4	.032719080665409	2579935404	= 3.141031743879264
192 ×	5	.016361730547870	1772725885	= 3.141452265191040
384 ×	6	.008181132952953	7982770512	= 3.141555053934336
768 ×	7	.004090603756546	7977365189	= 3.141583685027328
1536 ×	8	.002045306156319	2273950508	= 3.141590256105984
3072 ×	9	.001022653612915	7796943436	= 3.141591898874880
6144 ×	10	.000511326873302	4221834471	= 3.141592309567488
12288 ×	11	.000255663445004	-----	= 3.141592412233728
24576 ×	12	.000127831723547	8350493851	= 3.141592437891072
49152 ×	13	.000063915861904	4732978108	= 3.141592444305408
98304 ×	14	.000031957930970	0447026860	= 3.141592446074880
196608 ×	15	.000015978965486	3191307784	= 3.141592446201488
393216 ×	16	.000007989482743	4208153432	= 3.141592446271488
786432 ×	17	.0000039947413717	-----	= 3.1415924464287744
1572864 ×	18	.000001997370685866	-----	= 3.141592446453940224
3145728 ×	19	.000000998685342937	-----	= 3.141592446466523136

TABLE B

Side is to Side		As Loss is to Loss		
17	1	1	17	.0000000000548152861088220
17	2	2	17	.0000000000127399180702221
17	3	3	17	.0000000000008437784870086
17	4	4	17	.00000000000021131061251629
17	5	5	17	.0000000000002760761228788
17	6	6	17	.00000000000020444470896519
17	7	7	17	.00000000000008168799173397
17	8	8	17	.0000000000001164211782755
17	9	9	17	.00000000000019960171721350
17	10	10	17	.0000000000000540394788565
17	11	11	17	-----
17	12	12	17	.0000000000000267215932746
17	13	13	17	.0000000000000075727649685
17	14	14	17	.0000000000000003573711636
17	15	15	17	.0000000000000012765456446
17	16	16	17	.0000000000000008051066557
--	--	--	--	.0000000000758527031120600

FIRST THEOREM

Analytic Study of the Relation of the Radius to the Circumference

There is a true ratio existing between the radius and the quadrant of the circle, and it can be found as truly as the hypotenuse of a right angle triangle when the two sides are known. Fig. 1, Theorem First, and the following example worked out to four decimal places will fully explain.

I submit this proposition :

The square of the sum of the two sides AC and BC is to the difference of their squares (which is square of radius) as the sum of the two sides AC and BC is to their difference. Then :

To find the quadrant BEC add BD square + DC square - EF square to the square of the sum of the two sides AC and BC; for want of a better term, call it the square of the curve.




Fig. 1.

Solution of the proposition to four decimal places.

$$\begin{aligned} AC &= .50 \\ BC &= .7071 \end{aligned}$$

$$\begin{array}{r} 1.2071 \\ 1.2071 \\ \hline 12071 \\ 84497 \\ 24142 \\ 1.2071 \end{array}$$

1.45709041 square of the sum of the two sides AC and BC.

Find EF by taking $\frac{1}{2}$ the square of radius AB $= .5 \times .5 = \frac{.25}{2} = .125$

$$\begin{array}{r} \sqrt{.125} \quad .3535 \\ 9 \\ \hline 65 \quad 350 \\ 325 \\ \hline 703 \quad 2500 \\ 2109 \\ \hline 7065 \quad 39100 \\ 35325 \\ \hline \end{array}$$

$$\begin{array}{r}
\text{Then BD square + DC square} = .50 \\
- .3535 \\
\hline
.1465 = \text{EF} \\
.1465 \\
\hline
7325 \\
8790 \\
5860 \\
1465 \\
\hline
.02146225
\end{array}$$

$$\begin{array}{r}
\text{BD square + DC square} = .50 \\
- \text{EF square} = .02146225 \\
\hline
.47853775
\end{array}$$

$$\begin{array}{r}
\text{The square of the sum of the two other sides} = 1.45709041 \\
+ \text{BD square + DC square} - \text{EF square} = .47853775 \\
\hline
1.93562816
\end{array}$$

$$\begin{array}{r}
\sqrt{1.93562816} \text{ (1.3912 sum of the two sides.)} \\
1 \\
23 \overline{) 93} \\
69 \\
269 \overline{) 2456} \\
2421 \\
2781 \overline{) 3528} \\
2781 \\
\hline
74716
\end{array}$$

$$\begin{array}{r}
\text{The square of the sum of the two sides AC and BC} = 1.45709041 \\
+ \text{BD square + DC square} - \text{EF square} = .47853775 \\
\hline
1.93562816
\end{array}$$

$$\begin{array}{r}
\text{Then sq. of sum of 2 sides : sq. of radius :: sum of 2 sides : diff. of 2 sides} \\
1.93562816 \quad : \quad .25 \quad :: \quad 1.3912 \quad : \quad .1796
\end{array}$$

$$\begin{array}{r}
.25 \\
69560 \\
27824 \\
\hline
.347800000 \text{ (}.1796 \\
.193562816 \\
\hline
1542371840 \\
1354939712 \\
\hline
1874321280 \\
1742065344 \\
\hline
1322559360 \\
1161376896 \\
\hline
\text{Sum of the two sides, ... } 1.3912 \\
\text{Difference of the two sides, ... } .1796 \\
2 \overline{) 1.2116} \\
.6058 \\
.1796 \\
\hline
\text{Quadrant of circle, ... } .7854 \\
4 \\
\hline
3.1416
\end{array}$$

By calculating this proposition to fifteen decimal places the result will be 3.1419413724.

[illegible]

7

To make this solution of the question better understood, take the same example where 12 is the greater radius and 2 is the minor radius, as in Fig. 5.

AH=12	$\sqrt{148} \approx 12.165525$
AC= 2	1
$12 \times 12 = 144$	22 $\overline{) 48}$
$2 \times 2 = 4$	44
148	241 $\overline{) 400}$
	241
BC=Diagonal of the smaller radius.	2426 $\overline{) 15900}$
	14556
	24325 $\overline{) 134400}$
	121625
EF=One-half of difference between diagonal and smaller radius.	243305 $\overline{) 1277500}$
	1216525
	2433102 $\overline{) 6097500}$
	4866204
HO=The quadrant of the circumference.	2433104 $\overline{) 123129600}$
	12.165525=diagonal.
	2. =smaller radius.
	14.165525=sum of two sides.
	14.165525
	70827625
	28331050
	70827625
	70827625
	84993150
	14165525
	56662100
	14165525
	200.662098525625
	47.656855
	248.318953=sum of squares of two sides.
	$\sqrt{248.318953} \approx 15.758139$ =sum of the sides.
	1
	25 $\overline{) 148}$
	125
	307 $\overline{) 2331}$
	2149
	3145 $\overline{) 18289}$
	15725
	31508 $\overline{) 256453}$
	252064
	315161 $\overline{) 438900}$
	315161
	3151623 $\overline{) 12373900}$
	9454869
	31516269 $\overline{) 291903100}$

Find diagonal BC Fig. 5.

$$\begin{array}{l} AB=2 \times 2=4 \\ AC=2 \times 2=4 \end{array} \quad \sqrt{8} \text{ } 2.828427$$

$$\begin{array}{r} 8 \quad 48 \overline{) 400} \\ \underline{384} \\ 1600 \\ 1124 \\ \underline{47600} \\ 45184 \\ \underline{241600} \\ 226256 \\ \underline{1534400} \\ 1131364 \\ \underline{402036} \end{array}$$

Diagonal 2.828427
- Radius 2.

$$\begin{array}{r} \underline{.828427} \\ .828427 \\ \underline{5798989} \\ 1656854 \\ 3313708 \\ 6627416 \\ 1656854 \\ 6674162 \\ \underline{2 \text{ } .686291294329} \end{array}$$

.343145647164 = half the
square of difference of
diagonal and radius.

$$12 \times 2 = 24 \times 2 = 48.$$

$$\begin{array}{r} - .343145 \\ \underline{47.656855} \end{array}$$

$$\text{Then } 248.318953 : 144 :: 15.758139 : 9.138134$$

$$\begin{array}{r} 144 \\ \underline{63032556} \\ 63032556 \\ 15758139 \\ \underline{2269.172016} \\ 248.318953 \overline{) 2269.172016} \text{ } 9.138134 = \text{difference of two sides.} \\ \underline{2234870577} \end{array}$$

$$\begin{array}{r} 343014390 \\ 248318953 \\ \underline{946954370} \\ 744956859 \\ \underline{1142022110} \\ 993275812 \\ \underline{148746298} \end{array}$$

15.758139 = sum of two sides.
9.138134 = difference “

$$\begin{array}{r} 2 \overline{) 6.620005} \\ 3.310002 \\ \underline{9.138134} \end{array}$$

$\left. \begin{array}{l} 12.448136 = \text{quadrant} \\ \underline{4} \\ 49.792544 = \text{circumference} \end{array} \right\} \text{ and } \left\{ \begin{array}{l} \text{the ellipsis whose greater radius is 12} \\ \text{and whose minor radius is 2.} \end{array} \right.$

After calculating the area of the circle and its circumference according to First Theorem (Fig. 1), and also calculating the circumference of the ellipsis according to the same Theorem (Fig. 5), getting a little greater answer than is given in geometry, I made a careful calculation by approxi-




Fig. 7.

mation, taking a six-sided polygon inside of a circle whose diameter is 1 (Fig. 7), being divided into six segments, each and every side being equal to radius. Then $DC = .5 \times 6 = 3$, as per table of answers of subdivisions of sides calculated up to the sixteenth division $.000007989482743 \times 393216$ (number of sides) $= 3.141592446271488$.

The remainder vanished after the sixteenth subdivision.

As all areas having like sides are in proportion to the squares of such like sides, so would all losses or remainders be in proportion to the squares of such sides. Look on table of answers up to the nineteenth subdivision calculated to fifteen decimal places. (Table A.) To the right I have carried out ten places more, calling them remainders or losses.

Table B will show calculations of the same.

By taking the seventeenth subdivision for the first term, calculating all the losses from Table A in the remainder column, adding them up I find $.0000000000758527031120600$, as this is the loss in calculating one side, there being $786432 \times .0000000000758527031120600 =$

$.0000586539930294522099200$	Losses added.
3.141592446271488	Answer at sixteenth subdivision.
$3.14165110026451745220992$	Answer.

If I had taken the eighteenth subdivision for the first term, the sides would have been doubled and the sides would have been diminished almost one-half, which would have increased the losses almost four times, so I will multiply

$.00005865399302945220992$	by 4
$.00023461597211780883968$	
3.141592446271488	
3.141827062243605	

This answer is obtained on the theory of obtaining the result by approximation, and I give it as evidence of the correctness of Theorem First, Fig. 1.

SECOND THEOREM

I have found another solution of the same question, which I will endeavor to explain by calling attention to Fig. 6.

Taking a circle whose diameter is 1, construct a square AB touching the circle, each side of the square being equal to the diameter. Then construct a minor square within the circle marked GF Fig. 6.

Then $\sqrt{AB \times GF}$ = the quadrant of the circle. Multiply the quadrant of the circle CCCC by AB and the result obtained will be the area of the circle.




Fig. 6.

I find by inspection that six-sixteenths of 1 square (or outside square) = .375 square, which is a little too small by a little over .005. This difference is found by calculating the difference of the squares.

The question arises why the factors $\frac{2}{3}$ and $\frac{5}{8}$ are used to obtain the foregoing difference. A circle whose diameter is 1 will have a radius = .5, and the square of radius = .25. Suppose the same diameter 1 is taken, Let .375 = smaller radius and

“ .625 = greater radius.

Then $.375 \times .625 = .234375$ = square of mean radius.

Then $.25 - .234375 = .015625 = \frac{1}{4}$ BC square.

(Fig. 7) $\frac{.625}{4}$

Then the square of the radius of a perfect circle being .25, is to the square of the mean radius of the imperfect circle as the difference between the squares of the radius of the perfect circle and the imperfect circle (.234375) would be to the imperfect circle's difference, or ratio.

Then $.25 : .234375 :: .015625 : .0146484375$

$ \begin{array}{r} .015625 \\ \hline 1171875 \\ 468750 \\ 1406250 \\ 1171875 \\ 234375 \\ \hline .003662109375 \end{array} $	$ \begin{array}{r} .25 \mid .003662109375 \mid .0146484375 \\ \hline 25 \\ \hline 116 \\ 100 \\ \hline 162 \\ 150 \\ \hline 121 \\ 100 \\ \hline 210 \\ 200 \\ \hline 109 \\ 100 \\ \hline 93 \\ 75 \\ \hline 187 \\ 175 \\ \hline 125 \end{array} $
--	---

Then the mean between these two differences or ratios

$$.015625 \times .0146484375 = .0002288818359375$$

$$\text{Then } \sqrt{.0002288818359375} = .01512884119612272207$$

The mean square of the two ratios or difference

$$.01512884119612272207 \div 16 = .000945552574757670$$

add .0625

$$\begin{array}{r}
 .06344555257475767 \\
 \hline
 6
 \end{array}$$

.38067331544854602 Area of inside square
as in former example.

I calculated the six-sided polygon inside of a circle whose diameter is 1 to over three million sides, and by bringing up the losses I find a result of 3.1418 +. But the two foregoing Theorems or Propositions are not by Approximation and are either true or false. I believe them to be true and correct.

EBENEZER ERSKINE WALLACE.